

**Shahoon Hossein 1\***  
**Shahoon Shadi 2**  
**Sahabekhtiari Mehran 3**

1- Associate Prof. of Oral &  
Maxilofacial Surgery Group  
(Shahed U.M.S)

2- Student in Dentistry faculty

3- Dentist, Researcher

**\* Corresponding author:**

Shahed dental School west italia St, Vesal Ave. Tehran, Iran

Tel: 021-88681734

E-mail:

[Shahoonh@yahoo.com](mailto:Shahoonh@yahoo.com)

Journal of Medical Council of Islamic Republic of Iran, VOL. 32, NO. 1, Spring 2014: 88-94

● ORIGINAL ARTICLE CODE:06

## **Evaluation of permanent teeth extraction and related risk factors in patients of karaj central health city year in 91-92**

### **Abstract**

**Introduction:** Because of prevalence of permanent tooth extraction and it's related complication and due to bether alternative ways of tooth repair, this study is done in Health Centers of karaj province to evaluate the etiologies and related risk factors.

**Methods:** Survey was done by cross-sectional method on 240 patients. After extraction of data/factors like etiology of tooth remoral/ age/ sex/ educational level/ cigarette somking and History of systemic disease were analyzed by SPSS software version 10.

**Results:** In present study, the most common reason of tooth extraction are decaying (72.1%), periodontal disease (17.5%), prosthetic reasons (7.5%), orthodontics causes (1.7%) and other reason (1.3%).

(54.6%) were women and (45.4%) were men. In term of educational level (45.8%) were in undergratuate level and (8.3%) had university degree. Except sex, in all other factors p value was <0.05.

**Conclusion:** Because caries and periodontal diseases were the most causes of permanent tooth extraction in referred people development of the tooth and oral health system including an efficient program that educate people about prevention is needed.

**Keywords:** Extraction of teeth ,causes of extraction