

Morovvati Saeid 1*
Saghafinia Masoud 2
Amirpour Amraii Sara 3
Zahed Shekar Abi Hosna 3
Shahbazi Nastaran 3

1- Associate Prof. of Molecular Biology (Baq.U.M.S)

2- Associate Prof. of Anesthesiology, (Baq.U.M.S)

3- BSc in Cellular and Molecular Biology, Tehran Medical Unit, Islamic Azad University.

*** Corresponding author:**

Genetic Center, Baghiatolla University of Medical Sciences, Molla Sadra St. Tehran, Iran

Tel: 021-88039883

E-mail: morovvati@hotmail.com

Journal of Medical Council of Islamic Republic of Iran, VOL. 31, NO. 3, Autumn 2013: 265-271

CASE REPORT CODE:26

A report of two members of an Iranian family affected by autosomal recessive osteopetrosis

Abstract

Introduction: In the rare hereditary bone disorder of osteopetrosis, reduced bone resorption function leads to both the development of densely sclerotic fragile bones and progressive obliteration of the marrow spaces and cranial foramina. Marrow obliteration, typically associated with extramedullary hemopoiesis and hepatosplenomegaly, results in anemia and thrombocytopenia; and nerve entrapment accounts for progressive blindness and hearing loss. Severe infantile or malignant osteopetrosis is the worst type of the disease which has poor prognosis. In this study we report two cases of severe infantile or malignant type of the disease in an Iranian family.

Methods: Our two patients were children of a family that wife was the grandchild of husband's aunt. Two weeks after birth, the first patient had episodes of seizure and spastic in extremities. Gradually, the patient found upper and lower respiratory problems and horizontal nystagmus. X-Ray of the hand and foot showed widening and increased bone density. Physical examination showed hepatosplenomegaly and petechiae in extremities. The patient expired due to cardiopulmonary arrest. The same episodes seizure happened for the 2nd patient 2 weeks after birth. Gradually, Asymmetry between eyes and eventual blindness confirmed by ophthalmologist. Finally the patient expired because of severe pneumonia.

Conclusion: Autosomal recessive osteopetrosis has been reported in most ethnic groups although as the disease is very rare it is more frequently seen in ethnic groups where consanguinity is common. For the first time we report two cases of severe infantile or malignant type of the disease in an Iranian family.

Keywords: Osteopetrosis, Autosomal recessive, Consanguinity