Downloaded from jmciri.ir on 2025-11-17 ]

Rahimi Sharbaf Fatemeh 1
Rabiei Maryam 2\*
Darvish Narenjbon Soodabeh 2
Habibi Omid 3

1- Assistant Prof. of Obstetrics & Gynecology (T.U.M.S)

2- Specialis in Obstetrics & Gynecology (T.U.M.S)

3- Specialis in Dentistry, Researcher

## \* Corresponding author:

Women Hospital, Tehran University of Medical Sciences, Tehran, Iran

**Tel:** 021-88902960 **Fax:** 021-88915959

E-mail: rahimish@sina.tums.ac.ir

Journal of Medical Council of Islamic Republic of Iran, VOL. 31, NO. 3, Autumn 2013: 265-271

**ORIGINAL ARTICLE** CODE:24

Evaluation of relation betwean periodontal disease and preeclampsia in pregnant women

## **Abstract**

**Introduction:** Pre-eclampsia is a complication of pregnancy characterized by systemic vascular dysfunction and pathological changes in placental arteries. Chronic periodontal infections are etiological factor in vascular diseases. The aim was to determine the relation between maternal periodontal disease, and risk of pre-eclampsia.

**Methods:** A case-control study was carried out on 60 pre-eclampsia women 60 controls were matched based age, body mass index, educational level and parity. Periodontal examination was performed; and maternal periodontitis was determined with Williams probe (UNC-15).

**Results:** Average robing depth was  $4.1 \pm 0.46$  in pre-eclampsia group and  $3.2 \pm 0.56$  in normal group (P<0.01) and site with a probing depth >4mm was  $5.76 \pm 2.63$  in pre-eclampsia group and  $3.35 \pm 1.69$  in normal group (P=0.002).

**Conclusion:** There was appositive association between periodontal disease and risk of pre-eclampsia.

Keywords: Periodontal disease, pre-eclampsia, probing, clinical attachment loss (CAL)