

**Tousi Parviz 1\***

1- Professor of Dermatology group, (SB.U.M.S)

**\* Corresponding author:**

Dermatology Group Center, Qods SQ, Tajrish, Tehran, Iran.

**Telfax:** 021-22741507

E-mail:

[SRC@sbmu.ac.ir](mailto:SRC@sbmu.ac.ir)

Journal of Medical Council of Islamic Republic of Iran, VOL. 32, NO. 2, Summer 2014: 182-188

● **REVIEW ARTICLE** CODE:16

## **Androgenetic Alopecia**

### **Abstract**

Androgenetic alopecia is an extremely common disorder affecting 70% of men and 40% of women in their life. Men present with gradual thinning in the temporal areas, producing a reshaping of the anterior part of the hairline and progress to vertex

balding. Women with androgenetic alopecia usually present with diffuse thinning on the crown. Only 2 drugs (minoxidil and finasteride) currently have FDA-approved indications for treatment of androgenetic alopecia. Due to the frequency and the often significant impairment of life perceived by the affected patients, diagnosis and treatment of androgenetic alopecia is particularly important. This article briefly reviews the definition, diagnosis and treatment options of androgenetic alopecia.

**Keywords:** Androgenetic Alopecia, Minoxidil, Finasteride